DRY EYE REGIMEN



SUITHWEST PRIOR TO YOUR PRE-OPERATIVE APPOINTMENT

To prepare your eyes for your pre-operative testing, please begin the following regimen to BOTH EYES as soon as possible and continue until instructed otherwise.

Note: This must be started no later than <u>2 weeks</u> before your pre-op appointment. Do not use make-up or lotion on your face, or eyes the day of your pre-operative appointment as doing so can interfere with your pre-operative measurements.

Preservative Free Artificial Tears – One drop to both eyes 3 – 4 times daily or as directed

Examples: Refresh Mega-3, Refresh Plus, Retain MGD, Oasis Tears Plus (In Office \$20.00 +Tax)

Warm Compresses – Use moist heat with massage to the base of the eyelashes nightly

- ❖ Moist heat used to open plugged glands, improve blood flow, regain free flowing oils
- ❖ Moist Heat Mask (Oasis): Available at check out for your convenience (\$15.00 +Tax), OTC or online to alleviate the need to re-heat a washcloth; heat 20 seconds in the microwave (Patented Medibeads absorb water molecules from the air and release them as clean moist heat)

Lid Scrubs – Gently scrub eyelid margins at lash base

- Gentle pressure applied to clean the base of the lashes with Q-tip, cloth, or pad to remove bacteria, debris, and other inflammatory material. Use diluted baby shampoo
- ❖ Effective tea tree oil products: Oasis Lid and Lash cleansing pads (In Office \$30.00 +Tax), Blephadex, Mediviz, or Optase can be purchased online. Avenova spray (by prescription only)

Nighttime "PM" gel drops – Once nightly before bedtime, OTC product. Humidify air; avoid fans

- Gently pull down lower lid and squeeze a small amount into both eyes Examples: Refresh gel drops, Genteal PM, or Systane PM
- ❖ Use before bed. Will temporarily blur vision so use just before bed; wash residue from lases in the morning

<u>CONTACT LENSES:</u> If you wear contact lenses, they <u>will need to be out</u> for a certain amount of time prior to your pre-operative appointment and prior to your surgery.

- ❖ Soft Contact Lenses 2 weeks prior to your pre-operative appointment for measurements
- ❖ Hard Contact Lenses 6-8 weeks prior to your pre-operative appointment for measurements (Unless otherwise noted by the doctor)